

Lambeth and Southwark Sexually Transmitted Infections quick reference treatment guideline for primary care

Shared Lambeth and Southwark Council guidelines, with NHS Lambeth Clinical Commissioning Group (CCG) and NHS Southwark Clinical Commissioning Group (CCG) clinical recommendation.

Principles of Treatment

1. Please refer to the most up to date BNF and Summary of Product Characteristics for full drug monographs which include further dosing and interaction information. ALWAYS check for hypersensitivity/allergy.
2. This is a quick reference guide. Please refer to the Public Health England, National Institute for Health and Care Excellence (NICE) Summary of antimicrobial prescribing guidance – managing common infections February 2019 and relevant British Association for Sexual Health and HIV (BASHH) guidelines for further information.
3. This guideline is for uncomplicated cases. For complicated cases (e.g. treatment failure/recurrent episodes/clinician concern) consider discussing or referring to [Sexual and Reproductive Health \(SRH\) services](#).
4. This guideline is based on the best available evidence at the time of development. Its application must be modified by professional judgement, based on knowledge about individual patient co-morbidities, potential for drug interactions and involve patients in management decisions.
5. The majority of this guideline provides dose and duration of treatment for adults. Doses may need modification for age, weight and renal function.
6. If diarrhoea or vomiting occurs due to an antibiotic or the illness being treated, the efficacy of hormonal contraception may be impaired and additional precautions should be recommended. Also see NICE Clinical Knowledge Summaries: [Diarrhoea - antibiotic associated](#)
7. Sexually Transmitted Infections (STI) may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and Hepatitis C only need to be tested in [high risk groups](#).

Approved by the Southwark Medicines Management Committee and Lambeth Borough Prescribing Committee in June 2019

Review date: June 2021 (or sooner if evidence changes)

These guidelines have been developed by NHS Lambeth CCG, NHS Southwark CCG, Consultant in Sexual Health - Guy's and St Thomas' NHS Foundation Trust (GSTFT) and Lambeth Public Health. The guideline is based on the Public Health England and NICE Summary of antimicrobial prescribing guidance – managing common infections February 2019, the British Association for Sexual Health and HIV (BASHH) guidelines and input from a specialist in sexual health.

Please direct any comments or queries to Medicines Optimisation: NHS Southwark CCG (email: SOUCCG.Medicines-Optimisation@nhs.net, tel: 020 7525 3253), NHS Lambeth CCG (email: LAMCCG.medicinesoptimisation@nhs.net, tel: 020 3049 4197)

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
<p>Gonorrhoea</p> <p>PHE & NICE</p> <p>BASHH</p>	<p>Prerequisite (if unable to complete the following refer to SRH services):</p> <ul style="list-style-type: none"> ➤ Bacterial culture should be taken in addition to Nucleic Acid Amplification Test (NAAT) before treatment ➤ Partner notification should be pursued in all patients ➤ Test of cure is needed in all patients 			<p>Refer to SRH services for culture and treatment.</p>	<p>If persisting symptoms/signs, then culture at least 72 hours after treatment and look for other co-infection.</p> <p>If asymptomatic, test two weeks after treatment.</p> <p>Antibiotic resistance is now very high and this concerns all antibiotic used to treat Gonorrhoea.</p> <p>Take advice/refer to SRH services in treatment failure. SRH must report all treatment failures to Public Health England,</p>
	<p>Intramuscular (IM) ceftriaxone 1g Stat</p>	<p>If cannot provide 1st line treatment refer to SRH services for culture and treatment.</p>			
<p>Chlamydia</p> <p>PHE & NICE</p> <p>BASHH</p>	<p>Prerequisite (if unable to complete the following refer to SRH services):</p> <ul style="list-style-type: none"> ➤ Partner notification should be pursued in all patients. ➤ Treat partners or refer partners to SRH services. ➤ Test for reinfection at 3-6 months following treatment if under 25 years and consider if over 25years and at high risk of re-infection. 			<p>Due to lower cure rate in pregnancy, test of cure at least three weeks after end of treatment.</p> <p>Azithromycin is the most effective option in pregnancy or breastfeeding.</p> <p>Oral azithromycin 1g Stat then 500mg OD for 2 days (total 3 days treatment) (off-label use).</p>	<p>Opportunistically screen all aged 15-24 years.</p> <p>Risk factors for infection include age under 25 years, a new sexual partner or more than one sexual partner in the past year and lack of consistent condom use.</p> <p>Patients should be advised to avoid sexual intercourse (including oral sex) until they and their partner (s) have completed treatment (or wait seven days if treated with azithromycin).</p>
	<p>Oral doxycycline 100mg BD 7 days</p>	<p>Oral ofloxacin 200mg BD 7 days</p>			

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
<p>Epididymo-orchitis or Epididymitis</p> <p>BASHH</p> <p>PHE & NICE</p>	<p>Prerequisite (if unable to complete the following refer to SRH services):</p> <ul style="list-style-type: none"> ➤ If Gonorrhoea suspected take bacterial culture in addition to NAAT at presentation due to high rates of antibiotic resistance. ➤ If the patient is Gonorrhoea positive, perform test of cure: <ul style="list-style-type: none"> • If by culture, perform at least 72 hours after completion of treatment. • If by NAAT, perform 3 weeks after completing treatment. 				<p>If no improvement after 3 days, reassess diagnosis and therapy.</p> <p>Further follow-up is recommended at 2 weeks after the initiation of treatment to assess compliance with treatment, partner notification and improvement of symptoms.</p> <p>Usually due to Gram-negative enteric bacteria in men over 35 years with low risk of STI.</p> <p>Patients should be advised to abstain from sexual intercourse until they and their partner have completed treatment and follow up in those with confirmed and suspected sexually transmitted infection.</p>
	<p>If STI suspected: Intramuscular (IM) ceftriaxone 1g stat, plus oral doxycycline 100mg BD for 10-14 days</p> <p>If suspect urinary tract infection (UTI) treat as per Southwark and Lambeth Antimicrobial Guideline for Primary Care 2018.</p>	<p>If cannot provide 1st line treatment refer to SRH services for culture and treatment.</p>			
<p>Likely Non-Gonococcal Urethritis (NGU)</p> <p>BASHH</p>	<p>Prerequisite (if unable to complete the following refer to SRH services):</p> <ul style="list-style-type: none"> ➤ Urethral microscopy to rule out gonorrhoea and to diagnose urethritis ➤ Perform STI screening and culture before treatment. 				<p>If no improvement or recurrent NGU refer to SRH services.</p> <p>Patients should be advised to avoid sexual intercourse (including oral sex) until they and their partner (s) have completed treatment.</p>
	<p>Oral doxycycline 100mg BD for 7 days</p>	<p>Oral ofloxacin 200mg BD or 400mg OD for 7 days</p>			

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
<p>Pelvic Inflammatory Disease (PID)</p> <p>PHE & NICE</p> <p>BASHH</p>	<p>Refer woman and contacts to SRH services for cervical microscopy and treatment.</p> <p>If prefer to treat in GP practice see 2nd line option →</p>	<p>Prerequisite (if unable to complete the following refer to SRH services):</p> <ul style="list-style-type: none"> ➤ Cervical microscopy to rule out PID. ➤ Always test for gonorrhoea and chlamydia. 		<p>Refer to gynaecology.</p>	<p>Cervical microscopy is a sensitive test to rule out PID.</p> <p>If gonorrhoea likely (partner has it, sex abroad, severe symptoms), use regimen with ceftriaxone, as resistance to quinolones is high.</p> <p>Review within 3 days' of initiating treatment, if no improvement, review diagnosis and treatment, consider referral.</p> <p>Further review at end of treatment may be useful to check symptoms and compliance with all advice.</p> <p>Patients should be advised to avoid oral or genital intercourse until they, and their partner(s), have completed their treatment.</p>
<p>Trichomonas Vaginalis (TV)</p> <p>BASHH</p> <p>PHE & NICE</p>	<p>Oral metronidazole 400mg BD for 5 days</p>			<p>Oral metronidazole 400mg BD for 5 days</p> <p>AVOID 2g single dose metronidazole</p>	<p>Treat partner(s) or refer partner(s) to SRH services.</p> <p>Complete test of cure only if still symptomatic following treatment or if symptoms recur.</p> <p>If treatment fails (on-going discharge or repeat positive test at four weeks) refer to SRH services.</p> <p>Advise to abstain from sex for at least one week until patient and partner(s) have completed treatment and follow-up.</p>

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
<p>Genital Herpes</p> <p>BASHH</p> <p>PHE & NICE</p>	<p>Oral aciclovir 400 mg TDS for 5 days</p>		<p>Suppressive treatment (if more than six episodes per year): Oral aciclovir 400 mg BD. Discontinue after a maximum of 12 months to reassess recurrence frequency.</p>	<p>Seek SRH advice.</p>	<p>First episode: treat within five days while new lesions are still forming, or if systemic symptoms persist and refer to SRH services.</p> <p>Review after 5 days and continue treatment if new lesions still appearing and/or complex disease.</p> <p>Refer to SRH services in 2 to 3 weeks if symptoms are not improving.</p> <p>Recurrent: self-care if mild, or short course antiviral treatment (1st line agent) if five or less episodes per year, or suppressive therapy can be initiated if there are six or more episodes per year.</p> <p>Self-care: saline bathing, analgesia, Petroleum jelly or Topical anaesthetic agents, e.g. over the counter (OTC) 5% lidocaine ointment may be useful to apply especially prior to micturition. Discuss transmission.</p>
<p>Genital Warts</p> <p>BASHH</p> <p>CKS</p>	<p>Apply podophyllotoxin cream (0.15%) or solution (0.5%) twice daily for 3 days followed by 4 days of no application, for 4 cycles.</p> <p>If ineffective after 4 cycles (ie: 4 weeks) try a different method. Unlicensed for extra-genital (ie: anal) warts.</p>	<p>Apply imiquimod 5% cream 3 nights a week (usually Mon / Wed / Fri) and then wash off each morning, for up to 16 weeks.</p>	<p>Cryotherapy - repeat at weekly intervals for 4 weeks.</p> <p><u>OR</u></p> <p>Refer to SRH services.</p>	<p>Refer to SRH services</p>	<p>Review at the end of a treatment course to monitor response and assess the need for further therapy.</p> <p>Ensure that all female patients are on a robust method of contraception for the duration of treatment. See Summary of Product Characteristics (SPC) for selected preparation.</p>

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
<p>Molluscum Contagiosum</p> <p>BASHH</p>	<p>Apply podophyllotoxin 0.5% solution twice daily for 3 days followed by 4 days of no application. Cycles can be repeated if necessary, for up to 4 cycles (unlicensed use).</p>	<p>Apply imiquimod 5% cream 3 nights a week (usually Mon / Wed / Fri) and then wash off each morning, for up to 16 weeks (unlicensed).</p> <p style="text-align: center;"><u>OR</u></p> <p>Cryotherapy</p>	<p>No treatment for immunocompetent patients as can resolve naturally.</p>	<p>Cryotherapy and other destructive methods are safe. AVOID podophyllotoxin and Imiquimod.</p>	<p>Ensure that all female patients are on a robust method of contraception for the duration of treatment. See Summary of Product Characteristics (SPC) for selected preparation.</p>
<p>Bacterial Vaginosis (BV)</p> <p>BASHH</p> <p>PHE & NICE</p> <p>CKS</p> <p>continued on the next page</p>	<p>Oral metronidazole 400 mg BD for 5 days</p>		<p>Metronidazole 0.75% vaginal gel. 5g applicator PV at night for 5 nights</p> <p style="text-align: center;"><u>OR</u></p> <p>Clindamycin 2% vaginal cream 5g applicator PV at night, for 7 nights</p>	<p>Routine treatment of asymptomatic pregnant women not recommended</p> <p>Treat if symptomatic</p> <p>Women with additional risk factors for preterm birth may benefit from treatment before 20 weeks gestation</p> <p>1st line oral metronidazole 400 mg BD for 7 days.</p> <p>AVOID 2g single dose oral metronidazole</p> <p>Alternatives are:</p> <p>Metronidazole 0.75% vaginal gel. 5g applicator PV at night for 5 nights</p> <p style="text-align: center;"><u>OR</u></p> <p>Clindamycin 2% vaginal</p>	<p>Treating partners does not reduce relapse. A test of cure is not needed if symptoms resolve.</p> <p>Women with BV should be screened for Trichomonas Vaginalis if at risk of STI.</p> <p>For persistent BV in women with an intrauterine contraceptive device, consider removing the device and advising the use of an alternative form of contraception.</p> <p>Treatment is indicated for symptomatic women. If asymptomatic can opt not to treat.</p>

Infection	1st line agent	2nd line agent	Other alternatives	Pregnancy and Breastfeeding	Follow up and Comments
<p>STIs may co-exist therefore consider screening for other STIs if positive for one or more STIs. Screening should include: Chlamydia, Gonorrhoea, HIV, Syphilis (and Trichomonas Vaginalis if patients or their partner are Black African/Caribbean). Hepatitis B and C only need to be tested in high risk groups.</p>					
				cream 5g applicator PV at night for 7 nights	
<p>Vulvo-vaginal candidiasis</p> <p>PHE & NICE</p> <p>BASHH</p>	<p>Purchase OTC clotrimazole pessaries/ cream or fluconazole oral capsule, if non-recurrent (≤ 2 episodes in 6 months)</p> <p>For further information see:</p> <ul style="list-style-type: none"> • NHS Lambeth CCG: 'Self-care with over the counter products' leaflet • NHS Southwark CCG: 'Are you Self Care Aware?' leaflet • Self Care Forum website • NHS Choices website 	<p>Recurrent (>4 episodes per year):</p> <p>150mg oral fluconazole every 72 hours for three doses induction, followed by 150mg once a week for six months maintenance</p>		<p>Treat if symptomatic, no evidence that asymptomatic women need treatment.</p> <p>OTC clotrimazole 100mg pessary PV at night for 6 nights.</p> <p>AVOID oral antifungal treatment</p>	<p>Follow-up is unnecessary if symptoms resolve. Test of cure is unnecessary.</p> <p>There is no evidence to support treatment of asymptomatic male partners in either episodic or recurrent vulvo-vaginal Candidiasis.</p> <p>OTC topical antifungal creams may be used in addition to oral / vaginal treatment if there are vulval symptoms.</p> <p>All topical and oral azoles give over 80% cure.</p>
<p>Syphilis</p>	<p>Refer to SRH services</p>				
<p>Mycoplasma Genitalium</p>	<p>Refer to SRH services in GSTT only (King's do not test for Mycoplasma Genitalium)</p>				

Sexual and Reproductive Health (SRH) services contact details	
Guy's and St Thomas' NHS Foundation Trust	<p>If your patient has a positive diagnosis and requires additional testing, treatment or partner notification you can book an appointment at Burrell Street for the following day at https://sxt.org.uk/pl/E4PKNF To book the appointment you need to add the patients name, their mobile number and there is a box where you can add notes for the clinic.</p> <p>Referrals from health professionals can be sent to:</p> <ul style="list-style-type: none"> • Email: gst-tr.referralsrsh@nhs.net • Address: Sexual and reproductive health Business Support Team Burrell Street 4-6 Railway Arches Burrell Street London SE1 0UN https://www.guysandstthomas.nhs.uk/our-services/sexual-health/referrals.aspx <p>Telephone advice from senior clinician: 020 7188 6666</p>
King's College Hospital NHS Foundation Trust	<p>GP Referral form should be emailed to kch-tr.outpatientofficer@nhs.net.</p> <p>For emergency referrals, contact the department on Tel: 020 3299 5000 to be put through to the relevant person. Or bleep the HIV/Sexual Health on-call doctor via switchboard Tel: 020 3299 9000</p> <p>For general enquiries about walk-in clinics and appointments, contact the department on Tel:020 3299 5000 https://www.kch.nhs.uk/service/a-z/sexual-health</p>
Lewisham and Greenwich NHS Trust	<p>Main office for Sexual Health Tel: 0203 049 3516 http://www.kisp.org.uk/contact/</p>

References

1. Summary of antimicrobial prescribing guidance – managing common infections (February 2019), Public Health England, National Institute for Health and Care Excellence <https://www.nice.org.uk/Media/Default/About/what-we-do/NICE-guidance/antimicrobial%20guidance/summary-antimicrobial-prescribing-guidance.pdf> . Last accessed 09/05/2019
2. Sexually Transmitted Infections in Primary Care 2013, Royal College of General Practitioners <http://www.rcgp.org.uk/clinical-and-research/clinical-resources/sexually-transmitted-infections-in-primary-care.aspx>. Accessed 02/11/2016
3. UK National Guideline for the management of Bacterial Vaginosis, 2012, British Association for Sexual Health and HIV <https://www.bashhguidelines.org/media/1041/bv-2012.pdf> . Accessed 19/10/2016
4. Bacterial vaginosis, Clinical Knowledge Summaries, <https://cks.nice.org.uk/bacterial-vaginosis#!topicsummary> Accessed 29/11/2017
5. United Kingdom National Guideline on the Management of Vulvovaginal Candidiasis (2007), British Association for Sexual Health and HIV <https://www.bashhguidelines.org/media/1155/united-kingdom-national-guideline-on-the-management-of-vulvovaginal-candidiasis.pdf> Accessed 29/11/2017
6. UK national guideline for the management of infection with Chlamydia trachomatis <https://www.bashhguidelines.org/media/1045/chlamydia-2015.pdf>. Accessed 29/11/2017
7. British Association for Sexual Health and HIV update on the treatment of Chlamydia trachomatis (CT) infection September 2018, <https://www.bashhguidelines.org/media/1191/update-on-the-treatment-of-chlamydia-trachomatis-infection-final-16-9-18.pdf>, Last accessed 09/05/2019
8. 2010 United Kingdom national guideline for the management of epididymo-orchitis, British Association for Sexual Health and HIV, <https://www.bashhguidelines.org/media/1062/3546.pdf> . Accessed 29/11/2017
9. British Association for Sexual Health and HIV national guideline for the management of epididymo-orchitis (2018 draft) <https://www.bashhguidelines.org/media/1204/bashh-eo-guideline-consultation-version.pdf>. Accessed 25/04/2019
10. UK national guideline for the management of anogenital herpes 2014, British Association for Sexual Health and HIV, https://www.bashhguidelines.org/media/1019/hsv_2014-ijstda.pdf Accessed 29/11/2017
11. UK National Guidelines on the Management of Anogenital Warts 2015, British Association for Sexual Health and HIV, <https://www.bashhguidelines.org/media/1075/uk-national-guideline-on-warts-2015-final.pdf>. Accessed 29/11/2017
12. British Association for Sexual Health and HIV national guideline for the management of infection with Neisseria gonorrhoeae (2019), <https://www.bashhguidelines.org/media/1208/gc-2019.pdf> . Last accessed 09/05/2019
13. UK national guideline for the management of Genital Molluscum in adults 2014 British Association for Sexual Health and HIV, https://www.bashhguidelines.org/media/1055/mc_2014-ijstda.pdf. Accessed 02/11/2016

14. BASHH UK National Guideline on the management of non-gonococcal urethritis May 2017, British Association for Sexual Health and HIV https://www.bashhguidelines.org/media/1146/ngu-update-05_2017-final.pdf Accessed 15/11/2017
15. Update to the 2015 BASHH UK National Guideline on the management of non-gonococcal urethritis November 2018, <https://www.bashhguidelines.org/media/1199/ngu-bashh-update-2018.pdf>, Last accessed 09/05/2019
16. United Kingdom National Guideline on the Management of Trichomonas vaginalis 2014, British Association for Sexual Health and HIV, https://www.bashhguidelines.org/media/1042/tv_2014-ijstda.pdf Accessed 15/11/2017
17. Warts – anogenital Clinical Knowledge Summaries <https://cks.nice.org.uk/warts-anogenital> Accessed 15/11/2017
18. United Kingdom National Guideline for the Management of Pelvic Inflammatory Disease (2019 Interim Update), <https://www.bashhguidelines.org/media/1217/pid-update-2019.pdf> Last accessed 09/05/2019